



PRESCRIBED BOWEL PREP INSTRUCTIONS: MOVIPREP

Please follow these instructions, not the ones on the MOVIPREP box. This prep is time sensitive and requires you to follow the instructions exactly.

What you will need:

- Moviprep (picked up from your pharmacy)
- Over the counter- Simethicone anti-gas product (Gas-X, Mylanta Gas, Maalox Anti-gas, etc.) – you will only need 4 tablets
- Clear liquids (see examples below)

The day before your colonoscopy:

- Drink only clear liquids throughout the day starting in the morning (see examples below).
- No red or purple liquids.
- No solid foods or soft foods are allowed.
- No milk or dairy products are allowed.

Clear liquids are those you can “see through”. Examples include:

- Water
- Apple Juice
- White cranberry juice
- White grape juice
- Tea (no milk or cream)
- Clear broth or bouillon
- Ginger Ale
- Lemon Lime soda / Lemonade (no pulp)
- Kool-Aid or other fruit flavored drinks
- Jell-O
- Ice popsicles
- Black coffee (No creamer or milk)
- Hard candies
- Carbonated Soda
- Sports drink (Gatorade, Vitamin Water, Powerade, etc.)


If you are having difficulty drinking the prep:

- You can mix your prep mid-day and put in the refrigerator until you are ready to drink.
- Put your prep over ice or use a straw to drink it.
- Use ginger-ale, lemon/lime wedges, or any other clear liquid in between the prep to make it more enjoyable.
- Suck on hard candies (no red or purple).
- Take a break from drinking, walk around, and then resume drinking the prep.
- For discomfort while wiping, use wet wipes and apply Vaseline/Desitin to the area.*
- Remind yourself, you are doing this for your own health.
- If you are having problems completing your prep or you are still passing brown fluid after completing your prep, please contact the office: 480-563-5757.

The day before your colonoscopy: / /

- To mix your prep:
 - The morning prior to your procedure you can prepare the Moviprep.
 - Empty one (1) “Pouch A” and one (1) “Pouch B” into provided container.
 - Add luke warm water to the fill line indicated on the container and mix to dissolve.
 - Once dissolved, place in the refrigerator.

- 6:00 pm** you will begin drinking your prep
 - The Moviprep container is divided by 4 marks on the side.
 - Every 15 minutes, drink the solution down to the next mark (approximately 8 ounces) until the solution is gone.
 - Drink an additional 16 ounces of clear liquids for hydration. You may consume this liquid while drinking your prep.



- 9:00 pm** you will take 2 simethicone anti-gas tablets with 8 ounces of clear liquid
- 10:00pm** you will take 2 simethicone anti-gas tablets with 8 ounces of clear liquid

NOTE: You may want to mix your next dose and place in the refrigerator for the morning. You will follow the same mixing instructions as provided above.

The day of your colonoscopy / / at : am/pm

- Mix your prep if you have not done so already. Follow the mixing instructions provided above.
- Begin drinking the second dose of your prep
 - The Moviprep container is divided by 4 marks on the side.
 - Every 15 minutes, drink the solution down to the next mark (approximately 8 ounces) until the solution is gone.
 - Drink an additional 16 ounces of clear liquids for hydration. You may consume this liquid while drinking your prep.

NOTE: You must complete the entire prep to ensure the most effective cleansing.

DO NOT DRINK ANYTHING AFTER : am/pm

You must not drink anything or put anything in your mouth, including your prep, after the stated time. This includes water, gum, mints, and hard candies. Failure to follow these instructions will result in a cancellation of your procedure.